

On the Convergence Rate in the Uniform Ergodic Theorem*

Takeshi Yoshimoto

Department of Mathematics, Toyo University, Kawagoe, Saitama 350, Japan

E

y Elsevier - Publisher Connector

INTRODUCTION

Let X be a complex Banach space with norm $\delta_X(\cdot)$ and $B[X, X]$ be the Banach algebra of bounded linear transformations from X to X with norm $\pi_B(\cdot)$.

The purpose of this paper is to investigate the speed of uniform convergence for Cesàro averages of quasi-compact operators in $B[X, X]$. A fundamental result in this direction is the following theorem which generalizes Kryloff and Bogoliouboff's result for Markov operators to the case of quasi-compact operators on general Banach spaces.

THEOREM (Yosida and Kakutani [12]). *If $T \in B[X, X]$ is quasi-compact and power-bounded then there exists a compact projection $\Pi \in B[X, X]$ such that*

$$\pi_B \left[\frac{T + T^2 + \cdots + T^n}{n} - \Pi \right] = O\left(\frac{1}{n}\right).$$

The power-boundedness of quasi-compact operators enables us to say even more. Then the following two theorems constitute the substance of the present paper.

THEOREM 1. *Let $T \in B[X, X]$ be quasi-compact and $\pi_B[T^n] \leq K$ ($n = 0, 1, 2, \dots$) for some constant $K > 0$. Then for any $\alpha > 0$ there exists a*

* Supported in part by the Special Research Funds from Toyo University.

compact projection $\Pi_\alpha \in B[X, X]$ such that

$$\pi_B \left[(A_n^\alpha)^{-1} \sum_{k=0}^n A_{n-k}^{\alpha-1} T^k - \Pi_\alpha \right] = O \left(\frac{1}{n^\omega} \right), \quad \omega = \min(\alpha, 1),$$

where the A_k^α 's are the (C, α) -coefficients of order α .

THEOREM 2. Let $\{T(t): t \geq 0\}$ be a uniformly continuous semigroup in $B[X, X]$. Suppose that (i) $\sup_{t \geq 0} \pi_B[T(t)] = M < \infty$, and (ii) $T(1)$ is quasi-compact. Then for any $\alpha \geq 1$ there exists a compact projection $\Pi_\alpha \in B[X, X]$ such that

$$\pi_B \left[\frac{\alpha}{u^\alpha} \int_0^u (u-t)^{\alpha-1} T(t) dt - \Pi_\alpha \right] = O \left(\frac{1}{u} \right).$$

In Section 1 we prove these theorems. Some related topics are discussed in Section 2. Section 3 is devoted to giving some applications of the results to investigate the asymptotic ergodic behaviors of stationary Markov transition functions.

1. PROOFS OF THE THEOREMS

Given a general semigroup G in $B[X, X]$ which contains the identity operator I , we denote by $\text{CO}(G)$ the convex hull of G and by $\overline{\text{CO}}(G)$ its closure in the uniform operator topology. The orbit of $x \in X$ under $\text{CO}(G)$ is denoted by $\text{OR}(x)$ and its strong closure is denoted by $\overline{\text{OR}}(x)$. Recall that the (C, β) -coefficients for a real number $\beta > -1$ are defined by

$$A_0^\beta = 1,$$

$$A_k^\beta = \binom{k+\beta}{k} = \frac{(\beta+1)(\beta+2)\dots(\beta+k)}{k!}, \quad k = 1, 2, 3, \dots$$

Then it is well known that

$$A_n^\beta = \sum_{k=0}^n A_{n-k}^{\beta-1} \sim n^\beta / \Gamma(\beta+1).$$

Proof of Theorem 1. For notational convenience we write

$$C_n^\alpha[T] = (A_n^\alpha)^{-1} \sum_{k=0}^n A_{n-k}^{\alpha-1}, \quad n = 0, 1, 2, \dots$$

The case $0 < \alpha \leq 1$ has already been proved in [11] in a more general setting. Thus it suffices to prove only the case $\alpha > 1$. First we observe that $\{C_n^\alpha[T]\}$ ($n = 0, 1, 2, \dots$) is a system of almost uniformly invariant integrals

for the cyclic semigroup $G(T)$, where $G(T) = \{T^n: n = 0, 1, 2, \dots\}$. It is clear that

$$\sup_{n \geq 0} \pi_B[C_n^\alpha[T]] < \infty, \quad C_n^\alpha[T] \in B[X, X], \quad C_n^\alpha[T]x \in \overline{\text{OR}}(x)$$

for every $x \in X$ and all $n = 0, 1, 2, \dots$. Moreover, we have

$$\begin{aligned} C_n^\alpha[T](T - I) \\ = (A_n^\alpha)^{-1} \left\{ -A_n^{\alpha-1} + \sum_{k=0}^{n-1} (A_{n-k}^{\alpha-1} - A_{n-k-1}^{\alpha-1})T^{k+1} + T^{n+1} \right\} \end{aligned}$$

so that

$$\begin{aligned} \pi_B[C_n^\alpha[T](T - I)] &\leq (A_n^\alpha)^{-1} \{ A_n^{\alpha-1} + K(A_n^{\alpha-1} + 2) \} \\ &\leq n^{-1}M_d(\alpha), \quad n = 1, 2, 3, \dots, \end{aligned}$$

where $M_d(\alpha)$ is some positive constant depending only on α . From this it follows that $\pi_B[C_n^\alpha[T](T - I)] \rightarrow 0$ as $n \rightarrow \infty$. We can therefore apply Eberlein's uniform ergodic theorem [4] to assert that there exists a compact projection $\Pi_\alpha \in B[X, X]$ with $T\Pi_\alpha = \Pi_\alpha T = \Pi_\alpha$ and $\Pi_\alpha C_n^\alpha[T] = C_n^\alpha[T]\Pi_\alpha = \Pi_\alpha$ ($n = 0, 1, 2, \dots$), such that $\Pi_B[C_n^\alpha[T] - \Pi_\alpha] \rightarrow 0$ as $n \rightarrow \infty$. Thus by [9, Theorem 4.2] we have the decomposition $X = N(I - T) \oplus (I - T)X$ and $(I - T)X$ is closed, where $N(I - T)$ stands for the null space of the operator $I - T$. Using the open mapping theorem, there exists a constant $\Gamma > 0$ such that to each $x \in X$ corresponds a $u \in X$ with

$$x - \Pi_\alpha x = (I - T)u, \quad \delta_X(u) \leq \Gamma \cdot \delta_X(x - \Pi_\alpha x).$$

Hence we have

$$\begin{aligned} (C_n^\alpha[T] - \Pi_\alpha)x &= (C_n^\alpha[T] - \Pi_\alpha)(I - \Pi_\alpha)x \\ &= (C_n^\alpha[T] - \Pi_\alpha)(I - T)u = C_n^\alpha[T](I - T)u \end{aligned}$$

and so, for all $n = 1, 2, 3, \dots$

$$\begin{aligned} \delta_X((C_n^\alpha[T] - \Pi_\alpha)x) &= \delta_X(C_n^\alpha[T](I - T)u) \\ &\leq \pi_B[C_n^\alpha[T](T - I)]\delta_X(u) \\ &\leq n^{-1}\{\Gamma \cdot M_d(\alpha) \cdot \pi_B[I - \Pi_\alpha]\}\delta_X(x) \end{aligned}$$

as was to be shown, and the proof is complete.

Proof of Theorem 2. For brevity we write $G = \{T(t): t \geq 0\}$ and

$$C_u^\alpha[G] = \frac{\alpha}{u^\alpha} \int_0^u (u - t)^{\alpha-1} T(t) dt, \quad u > 0.$$

Since we have

$$C_u^\alpha[G] = uo - \lim_{n \rightarrow \infty} \left\{ \frac{\alpha}{u^\alpha} \int_{1/n}^{u-1/n} (u-t)^{\alpha-1} T(t) dt \right. \\ \left. - \frac{\alpha}{u^\alpha} \int_{1/n}^{u-1/n} (u-t)^{\alpha-1} dt \right\},$$

where the integrals can be approximated by suitable Riemann sums, we see that

$$\sup_{u>0} \pi_B[C_u^\alpha[G]] < \infty, \quad C_u^\alpha[G] \in \overline{\text{CO}}(G), \quad C_u^\alpha[G]x \in \overline{\text{OR}}(x)$$

for every $x \in X$ and all $u > 0$. Moreover, for $0 < t < u$

$$C_u^\alpha[G](T(t) - I) = \frac{\alpha}{u^\alpha} \left\{ \int_u^{u+t} (u+t-s)^{\alpha-1} T(s) ds \right. \\ \left. - \int_0^t (u-s)^{\alpha-1} T(s) ds \right. \\ \left. + \int_t^u [(u+t-s)^{\alpha-1} - (u-s)^{\alpha-1}] T(s) ds \right\}$$

and hence

$$\pi_B[C_u^\alpha[G](T(t) - I)] \leq \frac{\alpha M}{u^\alpha} [u^\alpha - (u-t)^\alpha + tu^{\alpha-1}] \\ = \alpha M \left[\frac{t}{u} + \left\{ 1 - \left(1 - \frac{t}{u} \right)^\alpha \right\} \right] \rightarrow 0 \quad \text{as } u \rightarrow \infty.$$

This shows that $\{C_u^\alpha[G]\} (u > 0)$ is a system of almost uniformly invariant integrals for G . Thus, making use of the Eberlein theorem, there exists a compact projection $\Pi_\alpha \in B[X, X]$ such that $\pi_B[C_u^\alpha[G] - \Pi_\alpha] \rightarrow 0$ as $u \rightarrow \infty$. Now we may and shall assume that $u \geq 1$. Let $n = [u]$ (the integer part of u) so that $u = n + r$ with $0 \leq r < 1$. Then

$$C_u^1[G] = nu^{-1}C_n^1[T(1)]C_1^1[G] + ru^{-1}T(n)C_r^1[G].$$

This gives $\Pi_1 = EC_1^1[G]$, where $E = uo\text{-}\lim_{n \rightarrow \infty} C_n^1[T(1)]$ and $\pi_B[C_n^1[T(1)] - E] = O(n^{-1})$ which follows from the Yosida-Kakutani

theorem. Therefore

$$C_u^1[G] - \Pi_1 = nu^{-1}\{C_n^1[T(1)] - E\}C_1^1[G] \\ + (nu^{-1} - 1)EC_1^1[G] + ru^{-1}T(n)C_r^1[G]$$

so that

$$\pi_B[C_u^1[G] - \Pi_1] \leq K(u^{-1}), \quad u \geq 1$$

for some constant $K > 0$ independent of u . On the other hand, if $\alpha > 1$ then

$$C_u^\alpha[G] = \frac{\Gamma(\alpha + 1)}{\Gamma(2)\Gamma(\alpha - 1)} u^{-\alpha} \int_0^u t(u-t)^{\alpha-2} C_t^1[G] dt, \\ u^{-\alpha} \int_0^u t(u-t)^{\alpha-2} dt = \frac{\Gamma(2)\Gamma(\alpha - 1)}{\Gamma(\alpha + 1)}.$$

Hence from the above estimates we have

$$\pi_B[C_u^\alpha[G] - \Pi_1] \leq \frac{\Gamma(\alpha + 1)}{\Gamma(2)\Gamma(\alpha - 1)} u^{-\alpha} \int_0^u t(u-t)^{\alpha-2} \\ \pi_B[C_t^1[G] - \Pi_1] dt \\ \leq \frac{\Gamma(\alpha + 1)}{(\alpha - 1)\Gamma(2)\Gamma(\alpha - 1)} \\ \times \left\{ K + \sup_{0 \leq t \leq 1} \pi_B[C_t^1[G] - \Pi_1] \right\} \cdot u^{-1} \\ = u^{-1}M_c(\alpha), \quad u \geq 1.$$

This implies that $\Pi_\alpha = \Pi_1$ for all $\alpha \geq 1$ and the proof is complete.

Remark. It is worthwhile to notice that the following statements can be proved by using the same method as above.

(1) If $T \in B[X, X]$ is quasi-compact and $T^n/n^\xi \rightarrow 0$ ($n \rightarrow \infty$) in the weak operator topology for some ξ with $0 < \xi < 1$, then for any $\alpha \geq 1$ there exists a compact projection $\Pi_\alpha^{(d)} \in B[X, X]$ such that

$$\pi_B[C_n^\alpha[T] - \Pi_\alpha^{(d)}] = O(n^{-(1-\xi)}).$$

(2) If $G = \{T(t): t \geq 0\}$ is a uniformly continuous semigroup with $T(1)$ quasi-compact and $T(t)/t^\xi \rightarrow 0$ ($t \rightarrow \infty$) in the uniform operator

topology for some ξ with $0 < \xi < 1$, then for any $\alpha \geq 1$ there exists a compact projection $\Pi_\alpha^{(c)} \in B[X, X]$ such that

$$\pi_B[C_u^\alpha[G] - \Pi_\alpha^{(c)}] = O(u^{-(1-\xi)}).$$

2. FURTHER DISCUSSIONS

Let $G = \{T(t): t \geq 0\} \subset B[X, X]$ be a uniformly continuous semigroup of type ω_0 :

$$\omega_0 = \inf_{t>0} t^{-1} \log \pi_B[T(t)] = \lim_{t \rightarrow \infty} t^{-1} \log \pi_B[T(t)].$$

The Abel mean for G is defined by

$$J_\lambda[G] = \int_0^\infty e^{-\lambda t} T(t) dt, \quad \lambda > \omega_0.$$

Then the infinitesimal generator A of G exists and is bounded, and the resolvent $R(\lambda; A)$ of A is identical with $J_\lambda[G]$ for $\lambda > \omega_0$. Furthermore

$$\lambda J_\lambda[G] = \frac{\lambda^{\alpha+1}}{\Gamma(\alpha+1)} \int_0^\infty t^\alpha e^{-\lambda t} C_t^\alpha[G] dt, \quad \alpha > 0, \lambda > \max(0, \omega_0).$$

THEOREM 3. *Suppose that for each $t > 0$, $T(t)$ is compact, with $T(t)/t \rightarrow 0$ as $t \rightarrow \infty$ in the uniform operator topology. Then:*

(1) *For each $\lambda > 0$ there exists a compact projection $E_\lambda \in B[X, X]$ such that*

$$\pi_B[C_n^1[\lambda J_\lambda[G]] - E_\lambda] = O\left(\frac{1}{n}\right),$$

$$\pi_B[C_n^1[\lambda R(\lambda; A)] - E_\lambda] = O\left(\frac{1}{n}\right).$$

(2) *There exists a compact projection $\Pi \in B[X, X]$ such that*

$$\pi_B[C_u^\alpha[G] - \Pi] \rightarrow 0 \quad \text{as } u \rightarrow \infty \ (\alpha \geq 1),$$

$$\pi_B[\lambda J_\lambda[G] - \Pi] \rightarrow 0 \quad \text{as } \lambda \rightarrow 0+.$$

(3) $X = N(I - \lambda J_\lambda[G]) \oplus (I - \lambda J_\lambda[G])X$ and $(I - \lambda J_\lambda[G])X$ is closed for each $\lambda > 0$.

(4) $X = N(A) \oplus R(A)$ and $R(A)$ (the range of A) is closed.

(5) The point 1 is a simple pole of $\lambda J_\lambda[G]$ for each $\lambda > 0$.

(6) The point 0 is a simple pole of $R(\lambda; A)$ with residue Π in (2).

Proof. If t is sufficiently large then by assumption, $\pi_B[T(t)] \leq t$ and so

$$\omega_0 = \lim_{t \rightarrow \infty} t^{-1} \log \pi_B[T(t)] \leq \lim_{t \rightarrow \infty} t^{-1} \log t = 0.$$

Let $\lambda > 0$ be fixed. We observe that $J_\lambda[G]$ is compact. To show this we write

$$J_\lambda^{(n)}[G] = \frac{1}{n} \sum_{k=1}^{\infty} e^{-\lambda k/n} T\left(\frac{k}{n}\right), \quad n = 1, 2, 3, \dots,$$

$$J_{\lambda, N}^{(n)}[G] = \frac{1}{n} \sum_{k=1}^N e^{-\lambda k/n} T\left(\frac{k}{n}\right), \quad n = 1, 2, 3, \dots; N = 1, 2, 3, \dots$$

Fix $\varepsilon > 0$ arbitrarily. Then there is a number $t_0 = t_0(\varepsilon) > 0$ and a constant $M > 0$ such that

$$\pi_B[T(t)] \leq \varepsilon \cdot t, \quad t > t_0$$

$$\pi_B[T(t)] \leq M, \quad 0 \leq t \leq t_0.$$

Put $N_0(\varepsilon, n) = [nt_0] + 1$ and $N_0 = N_0(\varepsilon, n)$. For each fixed n one gets

$$\begin{aligned} \pi_B[J_\lambda^{(n)}[G] - J_{\lambda, N}^{(n)}[G]] &\leq \varepsilon \cdot \sum_{k=N}^{\infty} e^{-\lambda k/n} \cdot \frac{k}{n} \\ &\leq \left\{ \frac{e^{-\lambda/n}}{n(1 - e^{-\lambda/n})} + \frac{ne^{-2\lambda/n}}{n^2(1 - e^{-\lambda/n})^2} \right\} \cdot \varepsilon, \quad N \geq N_0. \end{aligned}$$

Therefore $J_\lambda^{(n)}[G]$ is compact since $J_{\lambda, N}^{(n)}[G]$ is compact for all $N = 1, 2, 3, \dots$. Furthermore

$$\begin{aligned} J_\lambda[G] - J_\lambda^{(n)}[G] &= \sum_{k=1}^{\infty} \left\{ \int_{k/n}^{(k+1)/n} e^{-\lambda t} T(t) dt - \frac{1}{n} e^{-\lambda k/n} T\left(\frac{k}{n}\right) \right\} \\ &\quad + \int_0^{1/n} e^{-\lambda t} T(t) dt \\ &= \sum_{k=1}^{\infty} e^{-\lambda k/n} T\left(\frac{k}{n}\right) \left\{ \int_0^{1/n} (e^{-\lambda t} - 1) T(t) dt \right. \\ &\quad \left. + \int_0^{1/n} (T(t) - I) dt \right\} \\ &\quad + \int_0^{1/n} e^{-\lambda t} T(t) dt. \end{aligned}$$

However, we have for $n > 1/t_0$ the estimates

$$\begin{aligned}
 & \pi_B \left[\frac{1}{n} \sum_{k=1}^{\infty} e^{-\lambda k/n} T \left(\frac{k}{n} \right) \right] \\
 & \leq \frac{1}{n} \left\{ \sum_{k=1}^{N_0-1} e^{-\lambda k/n} \pi_B \left[T \left(\frac{k}{n} \right) \right] + \sum_{k=N_0}^{\infty} e^{-\lambda k/n} \pi_B \left[T \left(\frac{k}{n} \right) \right] \right\} \\
 & \leq \frac{1}{n} \left\{ M \cdot \sum_{k=1}^{\infty} e^{-\lambda k/n} + \varepsilon \cdot \sum_{k=1}^{\infty} e^{-\lambda k/n} \cdot \frac{k}{n} \right\} \\
 & = \frac{Me^{-\lambda/n}}{n(1 - e^{-\lambda/n})} + \frac{\varepsilon}{n} \left\{ \frac{e^{-\lambda/n}}{n(1 - e^{-\lambda/n})} + \frac{ne^{-2\lambda/n}}{n^2(1 - e^{-\lambda/n})^2} \right\}
 \end{aligned}$$

and

$$\pi_B \left[n \int_0^{1/n} (e^{-\lambda t} - 1) T(t) dt \right] \leq nM \int_0^{1/n} (1 - e^{-\lambda t}) dt.$$

Hence $\pi_B[J_\lambda^{(n)}[G] - J_\lambda[G]] \rightarrow 0$ as $n \rightarrow \infty$ and $J_\lambda[G]$ is compact. Using the equation

$$(J_\lambda[G])^n x = \frac{1}{(n-1)!} \int_0^\infty e^{-\lambda t} t^{n-1} T(t) x dt$$

we see by a direct computation that $\pi_B[(\lambda J_\lambda[G])^n/n] \rightarrow 0$ as $n \rightarrow \infty$ [8, Lemma 1]. Thus (1) follows at once from [7, Theorem 2.1] or [11, Theorem 3.2]. Noting that $\pi_B[C_n^1[J_1[G]] - E_1] \rightarrow 0$ as $n \rightarrow \infty$ taking $\lambda = 1$ in (1), we see by [8, Theorem] that there exists a compact projection $\Pi \in B[X, X]$ such that $\pi_B[C_u^1[G] - \Pi] \rightarrow 0$ as $u \rightarrow \infty$. From this we have for $\alpha > 1$

$$\begin{aligned}
 & \pi_B[C_u^\alpha[G] - \Pi] \\
 & \leq \frac{\Gamma(\alpha+1)}{\Gamma(2)\Gamma(\alpha-1)} u^{-\alpha} \int_0^u t(u-t)^{\alpha-2} \pi_B[C_t^1[G] - \Pi] dt \rightarrow 0
 \end{aligned}$$

as $u \rightarrow \infty$

as before, and (2) follows. Part (1) and the Dunford uniform ergodic theorem [2, Theorem 3.16] give (3) at once. Parts (4) and (6) result from (2) and the Hille–Phillips theorem [6, Theorem 18.8.4]. Part (5) is obtained from [3, Chap. VIII, Theorem 8.3] because of the compactness of $J_\lambda[G]$. Hence the proof of the theorem has been completed.

COROLLARY 1. *Let $G = \{T(t): t \geq 0\} \subset B[X, X]$ be a uniformly bounded and uniformly continuous semigroup. Suppose that $T(1)$ is quasi-compact. Then there exists a compact projection $\Pi \in B[X, X]$ such that*

$$\pi_B[\lambda J_\lambda[G] - \Pi] = O(\lambda) \quad (\lambda \rightarrow 0+).$$

THEOREM 4. *Let $\alpha \geq 1$ and let $\{T(t): t \geq 0\} \subset B[X, X]$ be a uniformly bounded strongly continuous semigroup with the infinitesimal generator A . Then for each $x \in X$ of the form $x = y + z$ with $y \in N(A)$ and $z \in R(A)$,*

$$\delta_X(C_u^\alpha[G]x - y) = O\left(\frac{1}{u}\right).$$

Proof. Let $z = Aw$. Then since $T(0) = I$ we have

$$\begin{aligned} C_u^1[G]x - y &= u^{-1} \int_0^u \frac{d}{ds} T(s)w \, ds \\ &= u^{-1}\{T(u) - I\}w, \quad u > 0 \end{aligned}$$

so that

$$\delta_X(C_u^1[G]x - y) \leq u^{-1} \cdot \delta_X(w) \cdot \left(2 \sup_{u \geq 0} \pi_B[T(u)]\right), \quad u > 0.$$

Therefore we have for $\alpha > 1$

$$\begin{aligned} &\delta_X(C_u^\alpha[G]x - y) \\ &\leq \frac{\Gamma(\alpha + 1)}{\Gamma(2)\Gamma(\alpha - 1)} u^{-\alpha} \int_0^u t(u - t)^{\alpha-2} \delta_X(C_t^1[G]x - y) \, dt \\ &\leq u^{-1} \cdot \frac{\Gamma(\alpha + 1)\delta_X(w)}{(\alpha - 1)\Gamma(2)\Gamma(\alpha - 1)} \left(2 \sup_{u \geq 0} \pi_B[T(u)]\right), \quad u > 0 \end{aligned}$$

proving the theorem.

Theorem 4 is the (C, α) ($\alpha \geq 1$)-generalization of Goldstein, Radin, and Showalter's result [5, Theorem 1] concerning the speed of pointwise strong convergence of the $(C, 1)$ -averages on $N(A) \oplus R(A)$.

COROLLARY 2. *Let $G = \{T(t): t \geq 0\} \subset B[X, X]$ be a uniformly bounded and uniformly continuous semigroup with the infinitesimal generator A . Then for each $x \in X$ of the form $x = y + z$ with $y \in N(A)$ and $z \in R(A)$*

$$\delta_X(\lambda J_\lambda[G]x - y) = O(\lambda) \quad (\lambda \rightarrow 0+).$$

3. APPLICATIONS

Let $\Omega = [0, 1]$ and $\mathcal{B}[\Omega]$ = the class of all Borel subsets of Ω . Let X be the complex Banach space of all complex valued σ -additive set functions $x(\cdot)$ defined for all $A \in \mathcal{B}[\Omega]$ with norm $\delta_X(x(\cdot))$ = total variation of $|x(A)|$ on Ω . We consider a stationary Markov transition function $P(t, \xi, A)$ ($t \geq 0$, $\xi \in \Omega$, $A \in \mathcal{B}[\Omega]$) which satisfies the conditions:

- (1) $P(t, \cdot, A)$ is a Borel function of ξ for fixed t, A ;
- (2) $P(t, \xi, \cdot)$ is a probability measure in A for fixed t, ξ ;
- (3) $P(0, \xi, A) = e_A(\xi)$, $e_A(\xi)$ denotes the indicator function of A ;
- (4) $P(s + t, \xi, A) = \int_{\Omega} P(t, \eta, A)P(s, \xi, d\eta)$ (the Chapman-Kolmogorov equation).

We define with the kernel $P(t, \xi, A)$

$$T(t)x(A) = \int_{\Omega} x(d\xi)P(t, \xi, A), \quad t \geq 0, A \in \mathcal{B}[\Omega], x \in X.$$

Then $\{T(t): t \geq 0\}$ is a semigroup of positive linear operators in $B[X, X]$ with $T(0) = I$ and $\pi_B[T(t)] \leq 1$ ($t \geq 0$). In addition, we impose the following condition upon the function $P(t, \xi, A)$:

- (5) $\lim_{t \rightarrow 0+} P(t, \xi, A) = e_A(\xi)$ uniformly in ξ, A .

Given any $\varepsilon > 0$ there exists by condition (5) a number $\delta = \delta(\varepsilon) > 0$ such that

$$|P(t, \xi, A) - e_A(\xi)| < \varepsilon, \quad 0 \leq t < \delta$$

uniformly in ξ and A . Thus we have for any $x \in X$

$$\delta_X([T(t) - I]x(\cdot)) \varepsilon \cdot \delta_X(x(\cdot))$$

from which it follows that $\{T(t): t \geq 0\}$ is uniformly continuous.

THEOREM 5. *Let $\alpha \geq 1$ and suppose that $T(1)$ is quasi-compact. Then there exists a transition probability function $Q^{(\alpha)}(t, \xi, A)$ ($t \geq 0$, $\xi \in \Omega$, $A \in \mathcal{B}[\Omega]$) such that*

$$(i) \quad \text{l.u.b.}_{\xi \in \Omega, A \in \mathcal{B}[\Omega]} \left| \frac{\alpha}{u^\alpha} \int_0^u (u-t)^{\alpha-1} P(t+s, \xi, A) dt - Q^{(\alpha)}(s, \xi, A) \right| \\ = O\left(\frac{1}{u}\right), \quad s \geq 0;$$

$$(ii) \quad \int_{\Omega} P(t, \xi, d\eta) Q^{(\alpha)}(s, \eta, A) = Q^{(\alpha)}(t+s, \xi, A), \quad s, t \geq 0.$$

Proof. Apply Theorem 2 to the semigroup $G = \{T(t): t \geq 0\}$ to obtain a compact projection $\Pi_\alpha \in B[X, X]$ such that $\pi_B[C_u^\alpha[G] - \Pi_\alpha] = O(u^{-1})$. Using Π_α we define

$$Q^{(\alpha)}(t, \xi, A) = \Pi_\alpha P(t, \xi, A), \quad t \geq 0, \xi \in \Omega, A \in \mathcal{B}[\Omega].$$

Then we have

$$\begin{aligned} \text{l.u.b.}_{\xi \in \Omega, A \in \mathcal{B}[\Omega]} & \left| \frac{\alpha}{u^\alpha} \int_0^u (u-t)^{\alpha-1} P(t+s, \xi, A) dt - Q^{(\alpha)}(s, \xi, A) \right| \\ & \leq \pi_B[C_u^\alpha[G] - \Pi_\alpha] \cdot \text{l.u.b.}_{\xi \in \Omega} \delta_X(P(s, \xi, \cdot)), \end{aligned}$$

and (i) follows. Clearly $Q^{(\alpha)}(t, \cdot, A)$ is a Borel function of ξ for fixed t, A , and $Q^{(\alpha)}(t, \xi, \cdot)$ is a probability measure in A for fixed t, ξ . As for (ii) we have by (i) and (4)

$$\begin{aligned} Q^{(\alpha)}(t+s, \xi, A) &= \lim_{u \rightarrow \infty} \frac{\alpha}{u^\alpha} \int_0^u (u-\omega)^{\alpha-1} P(\omega+t+s, \xi, A) d\omega \\ &= \lim_{u \rightarrow \infty} \frac{\alpha}{u^\alpha} \int_0^u (u-\omega)^{\alpha-1} d\omega \left[\int_\Omega P(\omega+s, \eta, A) P(t, \xi, d\eta) \right] \\ &= \int_\Omega P(t, \xi, d\eta) \left[\lim_{u \rightarrow \infty} \frac{\alpha}{u^\alpha} \int_0^u (u-\omega)^{\alpha-1} P(\omega+s, \eta, A) d\omega \right] \\ &= \int_\Omega P(t, \xi, d\eta) Q^{(\alpha)}(s, \eta, A), \end{aligned}$$

and the theorem is proved.

It should be noticed that if $P(1, \xi, A)$ satisfies the Doeblin (D)-condition then the operator $T(1)$ is quasi-compact by the Yosida-Kakutani theorem [12, Theorem 13].

Let W be the infinitesimal generator of $\{T(t): t \geq 0\}$. From Theorem 4 we have

THEOREM 6. *Let $\alpha \geq 1$. If $P(t_0, \xi_0, \cdot) \in N(W) \oplus R(W)$ for some t_0 and ξ_0 then*

$$\delta_X \left(\frac{\alpha}{u^\alpha} \int_0^u (u-t)^{\alpha-1} P(t+t_0, \xi_0, \cdot) dt - P_N(t_0, \xi_0, \cdot) \right) = O\left(\frac{1}{u}\right),$$

where $P(t_0, \xi_0, \cdot) = P_N(t_0, \xi_0, \cdot) + P_R(t_0, \xi_0, \cdot)$, $P_N(t_0, \xi_0, \cdot) \in N(W)$, $P_R(t_0, \xi_0, \cdot) \in R(W)$.

Remark. For each $\xi \in \Omega$, $P(0, \xi, \cdot) \in D(W)$ (the domain of W) and we have

$$-WP(0, \xi, A) = \lim_{t \rightarrow 0+} \frac{e_A(\xi) - P(t, \xi, A)}{t}.$$

Indeed, this follows from [1, Chap. VI, Theorems 2.2 and 2.3]. With $WP(0, \xi, A)$ define

$$q(\xi) = -WP(0, \xi, \{\xi\}), \quad \xi \in \Omega,$$

$$q(\xi, A) = WP(0, \xi, A), \quad \xi \in \Omega - A, A \in \mathcal{B}[\Omega].$$

Then $q(\xi, A) \leq q(\xi)$ for all $\xi \in \Omega$ and all $A \in \mathcal{B}[\Omega]$ with $A \subset \Omega - \{\xi\}$. More precisely, the pair $[q(\cdot), q(\cdot, \cdot)]$ constitutes a standard pair of q -functions in the sense of Doob (see [1, Chap. VI, Theorem 2.4]). Then $P(t, \xi, \cdot) \in D(W)$ for each $t > 0$ and each $\xi \in \Omega$, and

$$\begin{aligned} WP(t, \xi, A) &= -q(\xi)P(t, \xi, A) \\ &\quad + \int_{\Omega - \{\xi\}} P(t, \eta, A)q(\xi, d\eta), \quad A \in \mathcal{B}[\Omega]. \end{aligned}$$

To see this, we need the following representation (see [1, p. 269])

$$\begin{aligned} P(t, \xi, A) &= \int_0^t \left\{ \int_{\Omega - \{\xi\}} e^{-q(\xi)s} P(t-s, \eta, A)q(\xi, d\eta) \right\} ds \\ &\quad + e^{-q(\xi)t} e_A(\xi). \end{aligned}$$

Thus, keeping ξ fixed and differentiating with respect to t we have

$$\frac{\partial P(t, \xi, A)}{\partial t} = -q(\xi)P(t, \xi, A) + \int_{\Omega - \{\xi\}} P(t, \eta, A)q(\xi, d\eta)$$

for $t > 0$, so that $P(t, \xi, \cdot) \in D(W)$ and $WP(t, \xi, A) = \partial P(t, \xi, A)/\partial t$.

Next let $J_\alpha = \text{so-lim}_{u \rightarrow 0+} C_u^\alpha[G]$ for any real $\alpha > 0$ and $G = \{T(t): t \geq 0\}$. From the uniform continuity of G it follows that $P(s, \xi, \cdot) \in D(J_\alpha)$ for any $s \geq 0$ and all $\xi \in \Omega$ and that

$$\lim_{u \rightarrow 0+} \frac{\alpha}{u^\alpha} \int_0^u (u-t)^{\alpha-1} P(t+s, \xi, A) dt = J_\alpha P(s, \xi, A)$$

uniformly in ξ, A . If $P(s_0, \xi, A)$ satisfies the Doeblin (D)-condition for some $s_0 > 0$ then $T(s_0)$ is quasi-compact. In this case, the set $\{C_u^\alpha[G]P(s_0, \xi, A): u > 0\}$ contains a weakly convergent subsequence $\{C_{u_j}^\alpha[G]P(s_0, \xi, A)\}$ ($j = 1, 2, 3, \dots$) with $u_j \rightarrow 0+$ as $j \rightarrow \infty$ and by [10, Theorem 1] we have $D(J_\alpha) = X$, where $J_\alpha^2 = J_\alpha$ and $J_\alpha T(t) = T(t)J_\alpha = T(t)$ for all $t > 0$.

REFERENCES

1. J. L. Doob, "Stochastic Processes," Wiley, New York, 1953.
2. N. Dunford, Spectral theory. I. Convergence to projections, *Trans. Amer. Math. Soc.* **54** (1943), 185–217.
3. N. Dunford and J. T. Schwartz, "Linear Operators. I. General Theory," Interscience, New York, 1958.
4. W. F. Eberlein, Abstract ergodic theorems and weak almost periodic functions, *Trans. Amer. Math. Soc.* **67** (1949), 217–240.
5. J. A. Goldstein, C. Radin, and R. E. Showalter, Convergence rates of ergodic limits for semigroups and cosine functions, *Semigroup Forum* **16** (1978), 89–95.
6. E. Hille and R. S. Phillips, "Functional Analysis and Semigroups," Amer. Math. Soc. Colloq. Publ., Amer. Math. Soc., Providence, RI, 1957.
7. S. Ladouceur and M. Weber, Speed of convergence of the mean average operator for quasi-compact operators, to appear.
8. M. Lin, On the uniform ergodic theorem, II, *Proc. Amer. Math. Soc.* **46** (1974), 217–225.
9. T. Yoshimoto, Vector-valued Hausdorff summability methods and ergodic theorems, *Proc. Amer. Math. Soc.* **107** (1989), 915–926.
10. T. Yoshimoto, Local mean ergodic theorems in Banach spaces, *J. Math. Anal. Appl.* **151** (1990), 396–404.
11. T. Yoshimoto, On the speed of convergence in the (C, α) -uniform ergodic theorem for quasi-compact operators, *J. Math. Anal. Appl.* **176** (1993), 413–422.
12. K. Yosida and S. Kakutani, Operator-theoretical treatment of Markoff's process and mean ergodic theorem, *Ann. of Math.* **42** (1941), 188–228.